arXiv:cond-mat/0607118v2 [cond-mat.stat-mech] 20 Nov 2006

Rigorous results on the local equilibrium kinetics of a protein folding model

Marco Zamparo[1, *] and Alessandro Pelizzola[1, 2, †]

[1] *Dipartimento di Fisica and CNISM, Politecnico di Torino,*
c. Duca degli Abruzzi 24, Torino, Italy
[2] *INFN, Sezione di Torino*

Abstract

A local equilibrium approach for the kinetics of a simplified protein folding model, whose equilibrium thermodynamics is exactly solvable, was developed in [1]. Important properties of this approach are (i) the free energy decreases with time, (ii) the exact equilibrium is recovered in the infinite time limit, (iii) the equilibration rate is an upper bound of the exact one and (iv) computational complexity is polynomial in the number of variables. Moreover, (v) this method is equivalent to another approximate approach to the kinetics: the path probability method. In this paper we give detailed rigorous proofs for the above results.

---

[*] Electronic address: marco.zamparo@polito.it

[†] Electronic address: alessandro.pelizzola@polito.it

I. INTRODUCTION

The present paper contains details of the proofs of a few rigorous results for the local equilibrium approach to the kinetics of the Wako–Saitô–Muñoz–Eaton (WSME) model of protein folding, which was developed in [1]. This is a highly simplified model where one aims to describe the kinetics under the assumption that it is mainly determined by the structure of the native state, whose knowledge is assumed. It is a one–dimensional model, with long–range, many–body interactions, where a binary variable is associated to each peptide bond (the bond between consecutive aminoacids). Two aminoacids can interact only if they are in contact in the native state and all the peptide bonds between them are in the ordered state. Moreover an entropic cost is associated to each ordered bond.

A homogeneous version of the model was first introduced in 1978 by Wako and Saitô [2, 3], who solved exactly the equilibrium thermodynamics and studied the appearance of phase transitions in the thermodynamic limit. The full heterogeneous case was later considered by Muñoz, Eaton and coworkers [4, 5, 6], who introduced the single (double, triple) sequence approximations, i.e. they considered only configurations with at most one (two, three) stretches of consecutive ordered bonds, for both the equilibrium and the kinetics. The equilibrium problem has been subsequently studied in [7], with exact solutions for homogeneous β–hairpin and α–helix structures, mean field approximation and Monte Carlo simulations. The exact solution for the equilibrium in the full heterogeneous case was given in [8]. Moreover, in [9] it was shown that the equilibrium probability has an important factorization property, which implies the exactness of the cluster variation method (CVM) [10, 11, 12], a variational method for the study of lattice systems in statistical mechanics. Recently the model has found various applications in the study of the kinetics of real proteins [13, 14, 15, 16] and, interestingly enough, in a problem of strained epitaxy [18, 19, 20].

In [1] we have studied the kinetics of the WSME model in a master equation framework by means of a local equilibrium approach [21, 22, 23], by assuming that the probability distribution factors at any time in the same way as the equilibrium one. A few rigorous results about the local equilibrium kinetics were reported without a detailed proof, since the main purpose of that paper was to show the relevance of such an approach for the kinetics of protein folding. In the present paper we shall give detailed proofs of those results. Sec. II will be dedicated to a description of the model and our approach to the kinetics. In Sec.

III we will concentrate on the properties of the local equilibrium approximation. In Sec. III A we shall show that the free energy never increases with time, that is a feature of the exact solution which is preserved by our technique. This result will allow us to prove, in Sec. III B, that the exact equilibrium probability distribution is recovered in the long time limit. The study of the asymptotic behaviour will then lead us to define an approximate relaxation rate, that is an upper bound of the exact one, as it will be discussed in Sec. III C. In Sec. IV we will focus on a physically relevant example of kinetics, in order to show how the local equilibrium approximation reduces the complexity of the kinetic problem, from exponential to polynomial in the number of variables. Finally Sec. V will be devoted to the proof of the equivalence between our local equilibrium approach and the path probability method [24, 25, 26], the generalization of the CVM to the kinetics. It must be noticed that the only property of the WSME model which underlies the results of Secs. III and V is that the equilibrium problem can be solved exactly by means of the CVM, i.e. its equilibrium distribution factors in some ways. As a consequence, our conclusions are valid for any other model with the same property.

II. THE MODEL AND THE LOCAL EQUILIBRIUM KINETICS

The model describes a protein of $N+1$ aminoacids as a chain of N peptide bonds (connecting consecutive aminoacids) that can live in only two states (native and unfolded) and can interact only if they are in contact in the native structure and if all bonds in the chain between them are native. To each bond is associated a binary variable m_i, $i \in \{1, \dots, N\}$, with values $0, 1$ for unfolded and native state respectively. The effective free energy of the model (often improperly called Hamiltonian) reads

$$H_N(m) \doteq \sum_{i=1}^{N-1} \sum_{j=i+1}^{N} \epsilon_{i,j} \Delta_{i,j} \prod_{k=i}^{j} m_k - RT \sum_{i=1}^{N} q_i (1 - m_i) \qquad (1)$$

where R is the gas constant and T the absolute temperature. The first term assigns an energy $\epsilon_{i,j} < 0$ to the contact (defined as in [6, 8]) between bonds i and j if this takes place in the native structure ($\Delta_{i,j} = 1$ in this case and $\Delta_{i,j} = 0$ otherwise). The second term represents the entropic cost $q_i > 0$ of ordering bond i.

In order to solve exactly the equilibrium problem it has been found useful [8, 9] to map the one–dimensional WSME model onto a two–dimensional model through the introduction of the variables $x_{i,j} \doteq \prod_{k=i}^{j} m_k$ which satisfy the short–range constraints $x_{i,j} = x_{i,j-1}x_{i+1,j}$ for $1 \leq i < j \leq N$. These can be associated to the nodes of a triangular shaped portion Λ of a two–dimensional square lattice, defined by $\Lambda = \{(i,j) \in \mathbb{N}^2 : 1 \leq i \leq j \leq N\}$. Let $\mathcal{C}_\Lambda$ be the set of all configurations x on Λ that fulfil previous constraints and rewrite Eq. (1) (divided by RT and leaving apart an additive constant) in the form

$$H_\Lambda(x) = \sum_{(i,j)\in\Lambda} h_{i,j} x_{i,j}. \tag{2}$$

¿From now on we shall concentrate on the above equation, where the $h_{i,j}$'s can be temperature dependent, without referring to the original protein folding (or strained epitaxy) problem. We will denote by p^e_Λ the corresponding Boltzmann distribution and by Z_Λ the partition function:

$$p^e_\Lambda(x) = \frac{\exp[-H_\Lambda(x)]}{Z_\Lambda} \tag{3}$$

and

$$Z_\Lambda = \sum_{x\in\mathcal{C}_\Lambda} \exp[-H_\Lambda(x)]. \tag{4}$$

This distribution has been shown [9] to factor as

$$p^e_\Lambda(x) = \prod_{\alpha\in\mathcal{A}} [p^e_\alpha(x_\alpha)]^{a_\alpha}, \tag{5}$$

where $\mathcal{A}$ is a set of local clusters $\alpha \subset \Lambda$ made of all square plaquettes ($a_\alpha = 1$), the triangles lying on the diagonal boundary ($a_\alpha = 1$) and their intersections, that is internal nearest–neighbour pairs ($a_\alpha = -1$) and single nodes ($a_\alpha = 1$). It can be easily checked that the coefficients a_α are the Möbius numbers [11] for the set $\mathcal{A}$. For each cluster $\alpha \in \mathcal{A}$ we denote by x_α ($x_{\Lambda\setminus\alpha}$) the projection of x onto α ($\Lambda \setminus \alpha$), by $\mathcal{C}_\alpha$ the set of all configurations on α that are projections of configurations on Λ, and define the cluster equilibrium probability as the marginal distribution

$$p^e_\alpha(x_\alpha) \doteq \sum_{x_{\Lambda\setminus\alpha}} p^e_\Lambda(x). \tag{6}$$

Notice that by definition $x_{(i,j)} = x_{i,j}$. It is important to point out that cluster probabilities allow to reconstruct a probability on the whole lattice. Let $\mathcal{D}_\Lambda$ be the set of all cluster probabilities $p = \{p_\alpha\}_{\alpha\in\mathcal{A}}$ relative to $\mathcal{A}$ satisfying the compatibility conditions $p_\beta(x_\beta) = \sum_{x_{\alpha\setminus\beta}} p_\alpha(x_\alpha)$ for $\alpha, \beta \in \mathcal{A}$ and $\beta \subset \alpha$. Then, for $p \in \mathcal{D}_\Lambda$, consider the function $P_\Lambda[p] : \mathcal{C}_\Lambda \to$

$\mathbb{R}_+$ defined as

$$P_\Lambda[p](x) = \prod_{\alpha \in \mathcal{A}} [p_\alpha(x_\alpha)]^{a_\alpha}. \tag{7}$$

It can be checked that, due to the nature of the lattice and constraints, $P_\Lambda[p]$ is a probability on $\mathcal{C}_\Lambda$ and p_α is its marginal distribution relative to α [9].

As a consequence of Eq. (5), the equilibrium problem can be solved exactly [9] by means of the CVM. Since the Boltzmann distribution minimizes the free energy and factors, restricting the variational principle to distributions with the same property, i.e. of the form $P_\Lambda[p]$, one finds that $p^e \in \mathcal{D}_\Lambda$ is the minimum of the Kikuchi free energy

$$F_\Lambda[p] \doteq \sum_{\alpha \in \mathcal{A}} a_\alpha \sum_{x_\alpha} [\ln p_\alpha(x_\alpha) + h_\alpha(x_\alpha)] p_\alpha(x_\alpha) \tag{8}$$

with respect to $p \in \mathcal{D}_\Lambda$. Here h_α are defined by $h_\alpha(x_\alpha) = \sum_{(i,j)\in\alpha} h_{i,j} x_{i,j}$ and it follows that $H_\Lambda(x) = \sum_{\alpha \in \mathcal{A}} a_\alpha h_\alpha(x_\alpha)$. This variational approach has been used in [1] as the starting point for a very accurate treatment of the kinetics.

The kinetic problem has been stated in the framework of a master equation approach. Denoting by $W_\Lambda(x' \to x) \geq 0$ the transition probability per unit time from the state x' to $x \neq x'$, we have to solve

$$\frac{d}{dt} p^t_\Lambda(x) = \sum_{x' \in \mathcal{C}_\Lambda} W_\Lambda(x' \to x) p^t_\Lambda(x') \tag{9}$$

where from probability normalization it follows that $W_\Lambda(x \to x)$ has to be such that

$$\sum_{x \in \mathcal{C}_\Lambda} W_\Lambda(x' \to x) = 0. \tag{10}$$

It is known [27] that if the Boltzmann distribution is a stationary point for the equation, that is

$$\sum_{x' \in \mathcal{C}_\Lambda} W_\Lambda(x' \to x) p^e_\Lambda(x') = 0, \tag{11}$$

and W_Λ is irreducible, then the solution of the master equation converges to p^e_Λ in the long time limit. In the following we shall assume that these conditions hold.

In [1] we have studied the above problem by means of a local equilibrium approach [21, 22, 23], that is we have assumed that, provided the initial condition p^0_Λ factors according to Eq. (5) as the equilibrium probability, the solution p^t_Λ of the master equation factors in the same way at any subsequent time. We are therefore dealing with a kinetic problem in a restricted probability space. With this simplification the master equation yields for the cluster probabilities

$$\frac{d}{dt}p^t_\alpha(x_\alpha) = \sum_{x' \in \mathcal{C}_\Lambda} W_\alpha(x' \to x_\alpha) P_\Lambda[p^t](x'), \tag{12}$$

where

$$W_\alpha(x' \to x_\alpha) \doteq \sum_{x_{\Lambda \setminus \alpha}} W_\Lambda(x' \to x). \tag{13}$$

By taking marginals of Eq. (12) one can verify that $p^t \in \mathcal{D}_\Lambda$ if $p^0 \in \mathcal{D}_\Lambda$, i.e. the above evolution preserves the compatibility conditions between the cluster probabilities.

In addition, it has been shown in [9] (where explicit expressions are given) that the equilibrium cluster probabilities can be written as linear functions of the expectation values

$$\xi^e_{i,j} \doteq \sum_{x \in \mathcal{C}_\Lambda} x_{i,j} P_\Lambda[p^e](x) = \sum_{x_\alpha} x_{i,j} p^e_\alpha(x_\alpha) \tag{14}$$

for $(i,j) \in \Lambda$ and any α containing the node (i,j). By introducing in the same way the time–dependent variables $\xi_{i,j}(t)$ our kinetic problem will be turned into

$$\frac{d}{dt}\xi(t) = f(\xi(t)), \tag{15}$$

where, if $\xi = \{\xi_{i,j}\}_{(i,j)\in\Lambda}$ is the collection of the above expectation values corresponding to p, $f = \{f_{i,j}\}_{(i,j)\in\Lambda}$ is defined by

$$f_{i,j}(\xi) = \sum_{x \in \mathcal{C}_\Lambda} \sum_{x' \in \mathcal{C}_\Lambda} x_{i,j} W_\Lambda(x' \to x) P_\Lambda[p](x'). \tag{16}$$

f will be computed explicitly in Sec. IV for a given, physically relevant, choice of W_Λ. Notice that, in the framework of the WSME protein folding model, $\xi_{i,j}(t)$ is but the probability of the stretch from i to j being native at time t.

III. PROPERTIES OF THE LOCAL EQUILIBRIUM KINETICS

A. Time behaviour of the free energy

We start the study of the properties of the local equilibrium approach by showing that the free energy, which reduces to the Kikuchi free energy (8) due to the approximation which assumes that the probability factors at any time in the same way as the equilibrium one, is never increasing. This is a fundamental property which holds for an exact solution and is preserved by our approximation.

As customary [27] we add to the Kikuchi free energy the constant $\ln Z_\Lambda$, where Z_Λ is the partition function (4), and with a slight abuse of notation we indicate again by F_Λ this new function. Notice that, using Eqs. (7), (5), (3) and the relation $H_\Lambda(x) = \sum_{\alpha \in \mathcal{A}} a_\alpha h_\alpha(x_\alpha)$, one can find that for $p \in \mathcal{D}_\Lambda$

$$\begin{aligned}
\sum_{\alpha \in \mathcal{A}} a_\alpha \sum_{x_\alpha} h_\alpha(x_\alpha) p_\alpha(x_\alpha) + \ln Z_\Lambda &= \sum_{\alpha \in \mathcal{A}} a_\alpha \sum_{x \in \mathcal{C}_\Lambda} h_\alpha(x_\alpha) P_\Lambda[p](x) + \ln Z_\Lambda \\
&= \sum_{x \in \mathcal{C}_\Lambda} [H_\Lambda(x) + \ln Z_\Lambda] P_\Lambda[p](x) \\
&= -\sum_{x \in \mathcal{C}_\Lambda} \ln p^e_\Lambda(x) P_\Lambda[p](x) \\
&= -\sum_{\alpha \in \mathcal{A}} a_\alpha \sum_{x \in \mathcal{C}_\Lambda} \ln p^e_\alpha(x_\alpha) P_\Lambda[p](x) \\
&= -\sum_{\alpha \in \mathcal{A}} a_\alpha \sum_{x_\alpha} \ln p^e_\alpha(x_\alpha) p_\alpha(x_\alpha) \qquad (17)
\end{aligned}$$

and obtain thus

$$F_\Lambda[p] = \sum_{\alpha \in \mathcal{A}} a_\alpha \sum_{x_\alpha} p_\alpha(x_\alpha) \ln \frac{p_\alpha(x_\alpha)}{p^e_\alpha(x_\alpha)}. \qquad (18)$$

The function F_Λ, that we shall continue to call Kikuchi free energy, is not negative since $F_\Lambda[p] \geq F_\Lambda[p^e] = 0$, and in the following will play the role of a Lyapunov function [28]. We now study its time behaviour, mimicking the exact approach [27]. Our results can easily be rewritten for a discrete time formulation.

Proposition: If p^t is the solution of Eq. (12) with initial condition p^0, then a function $L_\Lambda : \mathcal{D}_\Lambda \to \mathbb{R}$ exists such that

i) $L_\Lambda[p] \leq 0 \quad \forall p \in \mathcal{D}_\Lambda$;

ii) $L_\Lambda[p] = 0 \iff p = p^e$;

iii) $\frac{d}{dt} F_\Lambda[p^t] = L_\Lambda[p^t]$.

Proof. For $p \in \mathcal{D}_\Lambda$ let

$$\psi_\Lambda[p](x) \doteq P_\Lambda[p](x) / p^e_\Lambda(x). \qquad (19)$$

The time derivative of the free energy can be written as

$$\frac{d}{dt} F_\Lambda[p^t] = \sum_{\alpha \in \mathcal{A}} a_\alpha \sum_{x_\alpha} \left[\ln \frac{p^t_\alpha(x_\alpha)}{p^e_\alpha(x_\alpha)} + 1 \right] \frac{d}{dt} p^t_\alpha(x_\alpha)$$

$$= \sum_{\alpha\in\mathcal{A}} a_\alpha \sum_{x_\alpha} \left[\ln \frac{p^t_\alpha(x_\alpha)}{p^e_\alpha(x_\alpha)}\right] \frac{d}{dt} p^t_\alpha(x_\alpha), \tag{20}$$

where the last step follows from normalization of cluster probabilities. Using Eqs. (12), (13) and (19) respectively we have

$$\begin{aligned}
\frac{d}{dt} F_\Lambda[p^t] &= \sum_{\alpha\in\mathcal{A}} a_\alpha \sum_{x_\alpha} \left[\ln \frac{p^t_\alpha(x_\alpha)}{p^e_\alpha(x_\alpha)}\right] \sum_{x'\in\mathcal{C}_\Lambda} W_\alpha(x' \to x_\alpha) p^e_\Lambda(x') \psi_\Lambda[p^t](x') \\
&= \sum_{x\in\mathcal{C}_\Lambda} \sum_{x'\in\mathcal{C}_\Lambda} \left[\sum_{\alpha\in\mathcal{A}} a_\alpha \ln \frac{p^t_\alpha(x_\alpha)}{p^e_\alpha(x_\alpha)}\right] W_\Lambda(x' \to x) p^e_\Lambda(x') \psi_\Lambda[p^t](x') \\
&= \sum_{x\in\mathcal{C}_\Lambda} \sum_{x'\in\mathcal{C}_\Lambda} W_\Lambda(x' \to x) p^e_\Lambda(x') \psi_\Lambda[p^t](x') \ln \psi_\Lambda[p^t](x), \qquad (21)
\end{aligned}$$

and statement *iii*) of our proposition is obtained by setting

$$L_\Lambda[p] \doteq \sum_{x\in\mathcal{C}_\Lambda} \sum_{x'\in\mathcal{C}_\Lambda} W_\Lambda(x' \to x) p^e_\Lambda(x') \psi_\Lambda[p](x') \ln \psi_\Lambda[p](x). \tag{22}$$

Statements *i*) and *ii*) follow from the equivalence (to be proven later) of the above definition and

$$L_\Lambda[p] = \sum_{x\in\mathcal{C}_\Lambda} \sum_{x'\in\mathcal{C}_\Lambda\setminus\{x\}} W_\Lambda(x' \to x) p^e_\Lambda(x') \left[\psi_\Lambda[p](x') - \psi_\Lambda[p](x) + \psi_\Lambda[p](x') \ln \frac{\psi_\Lambda[p](x)}{\psi_\Lambda[p](x')}\right]. \tag{23}$$

The inequality $a - b + a \ln \frac{b}{a} \leq 0$ ($a, b \in \mathbb{R}_+$) implies *i*). In order to obtain *ii*) observe that the previous inequality reduces to an equality if and only if $a = b$ and then, by the irreducibility of W_Λ, $L_\Lambda[p]$ vanishes if and only if $\psi_\Lambda[p](x)$ is independent of x. $P_\Lambda[p] = p^e_\Lambda$ follows by normalization and $p = p^e$ by computing the marginal distributions of each member.

We complete our proof by showing the equivalence between Eq. (22) and (23). From now on let ψ be a positive function defined on $\mathcal{C}_\Lambda$. Notice first that, thanks to Eqs. (10) and (11), we have

$$\begin{aligned}
0 &= \sum_{x'\in\mathcal{C}_\Lambda} \left[\sum_{x\in\mathcal{C}_\Lambda} W_\Lambda(x' \to x)\right] p^e_\Lambda(x')\psi(x') - \sum_{x\in\mathcal{C}_\Lambda} \left[\sum_{x'\in\mathcal{C}_\Lambda} W_\Lambda(x' \to x) p^e_\Lambda(x')\right] \psi(x) \\
&= \sum_{x\in\mathcal{C}_\Lambda} \sum_{x'\in\mathcal{C}_\Lambda\setminus\{x\}} W_\Lambda(x' \to x) p^e_\Lambda(x')\psi(x') + \sum_{x\in\mathcal{C}_\Lambda} W_\Lambda(x \to x) p^e_\Lambda(x)\psi(x) + \\
&- \sum_{x\in\mathcal{C}_\Lambda} \sum_{x'\in\mathcal{C}_\Lambda\setminus\{x\}} W_\Lambda(x' \to x) p^e_\Lambda(x')\psi(x) - \sum_{x\in\mathcal{C}_\Lambda} W_\Lambda(x \to x) p^e_\Lambda(x)\psi(x) \\
&= \sum_{x\in\mathcal{C}_\Lambda} \sum_{x'\in\mathcal{C}_\Lambda\setminus\{x\}} W_\Lambda(x' \to x) p^e_\Lambda(x') \left[\psi(x') - \psi(x)\right]. \qquad (24)
\end{aligned}$$

In addition, using again Eq. (10), we can see that

$$\begin{aligned}
&\sum_{x\in\mathcal{C}_\Lambda}\sum_{x'\in\mathcal{C}_\Lambda} W_\Lambda(x'\to x)p^e_\Lambda(x')\psi(x')\ln\psi(x) \\
&= \sum_{x\in\mathcal{C}_\Lambda}\sum_{x'\in\mathcal{C}_\Lambda} W_\Lambda(x'\to x)p^e_\Lambda(x')\psi(x')\ln\psi(x) - \sum_{x'\in\mathcal{C}_\Lambda}\left[\sum_{x\in\mathcal{C}_\Lambda} W_\Lambda(x'\to x)\right]p^e_\Lambda(x')\psi(x')\ln\psi(x') \\
&= \sum_{x\in\mathcal{C}_\Lambda}\sum_{x'\in\mathcal{C}_\Lambda\setminus\{x\}} W_\Lambda(x'\to x)p^e_\Lambda(x')\psi(x')\ln\psi(x) + \sum_{x\in\mathcal{C}_\Lambda} W_\Lambda(x\to x)p^e_\Lambda(x)\psi(x)\ln\psi(x) + \\
&- \sum_{x\in\mathcal{C}_\Lambda}\sum_{x'\in\mathcal{C}_\Lambda\setminus\{x\}} W_\Lambda(x'\to x)p^e_\Lambda(x')\psi(x')\ln\psi(x') - \sum_{x\in\mathcal{C}_\Lambda} W_\Lambda(x\to x)p^e_\Lambda(x)\psi(x)\ln\psi(x) + \\
&= \sum_{x\in\mathcal{C}_\Lambda}\sum_{x'\in\mathcal{C}_\Lambda\setminus\{x\}} W_\Lambda(x'\to x)p^e_\Lambda(x')\psi(x')\ln\frac{\psi(x)}{\psi(x')}. \qquad (25)
\end{aligned}$$

B. Exactness of the equilibrium state

In spite of an approximate picture for nonequilibrium states, the local equilibrium approach allows to recover the exact probability distribution in the long time limit. This is due to the role of Lyapunov function [28] played by the Kikuchi free energy F_Λ, as stated in the following proposition.

Proposition: If p^t is the solution of Eq. (12) with initial condition p^0, then

$$\lim_{t\to+\infty} p^t = p^e. \qquad (26)$$

Proof. p^e is the stationary point of Eq. (12), as a consequence of Eqs. (5) and (11). We already know that a non–positive function L_Λ exists, such that $L_\Lambda[p]=0$ if and only if $p=p^e$ and $L_\Lambda[p^t]$ is the derivative of the free energy $F_\Lambda[p^t]$ with respect to time. Moreover p^e is the minimum of F_Λ. Then a theorem by Lyapunov [28] ensures that p^e is asymptotically stable.

C. Asymptotic behaviour

In the present section we shall discuss how the local equilibrium kinetics approaches the exact equilibrium state. This will lead us to define a relaxation rate for our approximation, and we shall show that this rate is an upper bound of the exact one.

The study of the asymptotic behaviour of the approximate evolution involves the linearized form of Eq. (12), for which, as we shall prove, some properties of the exact kinetic equations are still valid. To begin with we recall the algebraic structure underlying the near–equilibrium kinetics in the exact case.

Let us consider, on the vector space V_Λ of all real functions on $\mathcal{C}_\Lambda$, the scalar product

$$(\phi|\psi) \doteq \sum_{x\in\mathcal{C}_\Lambda} \frac{\phi(x)\psi(x)}{p^e_\Lambda(x)}. \tag{27}$$

It is known [27] that if the transition probability satisfies the detailed balance principle, that is

$$W_\Lambda(x' \to x)p^e_\Lambda(x') = W_\Lambda(x \to x')p^e_\Lambda(x), \tag{28}$$

then the linear operator $\mathbb{W}_\Lambda : V_\Lambda \to V_\Lambda$ defined by

$$(\mathbb{W}_\Lambda\phi)(x) = \sum_{x'\in\mathcal{C}_\Lambda} W_\Lambda(x' \to x)\phi(x') \tag{29}$$

is self-adjoint and negative semi-definite with respect to the above scalar product. Moreover, if W_Λ is irreducible then $(\phi|\mathbb{W}_\Lambda\phi) < 0$ provided that ϕ is not proportional to p^e_Λ, and the exact relaxation rate k^{ex} is given by

$$k^{\mathrm{ex}} = -\max\left\{\frac{(\phi|\mathbb{W}_\Lambda\phi)}{(\phi|\phi)} : \phi \in V_\Lambda \setminus \{0\} \text{ and } (\phi|p^e_\Lambda) = 0\right\}. \tag{30}$$

Let us move now to our approach and attempt to extend these results. From now on we shall assume detailed balance (28). Let us intoduce the vector space $\mathcal{V}_\Lambda$ of collections $u = \{u_\alpha\}_{\alpha\in\mathcal{A}}$ of real functions on $\mathcal{C}_\alpha$, for all $\alpha \in \mathcal{A}$, satisfying the conditions $\sum_{x_{\alpha\setminus\beta}} u_\alpha(x_\alpha) = u_\beta(x_\beta)$ if $\beta \subset \alpha$ and $\sum_{x_\alpha} u_\alpha(x_\alpha) = 0$. Notice that, since p^e_α is strictly positive and thanks to the previous conditions, for $u \in \mathcal{V}_\Lambda$ it is possible to find $\epsilon_0 > 0$ such that $p^e + \epsilon u \in \mathcal{D}_\Lambda$ if $\epsilon \in (-\epsilon_0, \epsilon_0)$. On the space $\mathcal{V}_\Lambda$ we can define a scalar product through the bilinear form

$$\langle u|v\rangle \doteq \sum_{\alpha\in\mathcal{A}} a_\alpha \sum_{x_\alpha} \frac{u_\alpha(x_\alpha)v_\alpha(x_\alpha)}{p^e_\alpha(x_\alpha)}. \tag{31}$$

It can be checked that $\langle u|u\rangle > 0$ for $u \in \mathcal{V}_\Lambda \setminus \{0\}$. Indeed, for $u \neq 0$,

$$\langle u|u\rangle = \left.\frac{d^2}{d\epsilon^2}F_\Lambda[p^e + \epsilon u]\right|_{\epsilon=0} \tag{32}$$

as it can be easily verified, and the result follows by the fact that p^e is the minimun of the free energy F_Λ. Let $\|u\|$ be the norm generated by this scalar product, i.e. $\|u\| = \sqrt{\langle u|u\rangle}$. Observe that the application $U_\Lambda : \mathcal{V}_\Lambda \to V_\Lambda$, defined by

$$U_\Lambda[u](x) = \left.\frac{d}{d\epsilon}P_\Lambda[p^e + \epsilon u](x)\right|_{\epsilon=0} = p^e_\Lambda(x)\sum_{\alpha\in\mathcal{A}} a_\alpha \frac{u_\alpha(x_\alpha)}{p^e_\alpha(x_\alpha)}, \tag{33}$$

preserves the scalar products, since

$$\begin{aligned}(U_\Lambda[u]|U_\Lambda[v]) &= \sum_{x\in\mathcal{C}_\Lambda}\sum_{\alpha\in\mathcal{A}} a_\alpha \frac{u_\alpha(x_\alpha)}{p^e_\alpha(x_\alpha)} U_\Lambda[v](x) \\ &= \frac{d}{d\epsilon}\sum_{\alpha\in\mathcal{A}} a_\alpha \sum_{x\in\mathcal{C}_\Lambda} \frac{u_\alpha(x_\alpha)}{p^e_\alpha(x_\alpha)} P_\Lambda[p^e+\epsilon v](x)\Bigg|_{\epsilon=0} \\ &= \frac{d}{d\epsilon}\sum_{\alpha\in\mathcal{A}} a_\alpha \sum_{x_\alpha} \frac{u_\alpha(x_\alpha)}{p^e_\alpha(x_\alpha)} [p^e_\alpha(x_\alpha)+\epsilon v_\alpha(x_\alpha)]\Bigg|_{\epsilon=0} \\ &= \sum_{\alpha\in\mathcal{A}} a_\alpha \sum_{x_\alpha} \frac{u_\alpha(x_\alpha)v_\alpha(x_\alpha)}{p^e_\alpha(x_\alpha)} = \langle u|v\rangle. \end{aligned} \tag{34}$$

The local equilibrium relaxation rate appears naturally through a linearization of the near–equilibrium kinetics. To deal with an evolution in the space $\mathcal{V}_\Lambda$, we set $p^t = p^e + u^t$, such that $u^t \in \mathcal{V}_\Lambda$ for all $t \in \mathbb{R}_+$. Then we substitute Eq. (12) with the equivalent one

$$\frac{d}{dt}u^t = \mathbb{T}_\Lambda u^t + R_\Lambda(u^t), \tag{35}$$

where $\mathbb{T}_\Lambda : \mathcal{V}_\Lambda \to \mathcal{V}_\Lambda$ is the linear operator defined by

$$\begin{aligned}(\mathbb{T}_\Lambda u)_\alpha(x_\alpha) &= \frac{d}{d\epsilon}\sum_{x'\in\mathcal{C}_\Lambda} W_\alpha(x'\to x_\alpha) P_\Lambda[p^e+\epsilon u](x')\Bigg|_{\epsilon=0} \\ &= \sum_{x'\in\mathcal{C}_\Lambda} W_\alpha(x'\to x_\alpha) U_\Lambda[u](x'). \end{aligned} \tag{36}$$

$\mathbb{T}_\Lambda$ is the operator corresponding to the Jacobian of the r.h.s. of Eq. (12) evaluated at $p = p^e$ and, as a consequence, there exists $\mathcal{I}_0 \subset \mathcal{V}_\Lambda$, neighborhood of 0, and a constant $M > 0$ such that

$$\|R_\Lambda(u)\| \le M\|u\|^2 \tag{37}$$

for all $u \in \mathcal{I}_0$.

$\mathbb{T}_\Lambda$ inherits its properties from $\mathbb{W}_\Lambda$ thanks to the relation

$$\langle u|\mathbb{T}_\Lambda v\rangle = (U_\Lambda[u]|\mathbb{W}_\Lambda U_\Lambda[v]), \tag{38}$$

which can be immediately verified. It follows that $\mathbb{T}_\Lambda$ is self-adjoint and negative definite. The first property is obvious, while for the second one it is sufficient to observe that

$$\langle u|\mathbb{T}_\Lambda u\rangle = (U_\Lambda[u]|\mathbb{W}_\Lambda U_\Lambda[u]) \le -k^{\text{ex}}(U_\Lambda[u]|U_\Lambda[u]) = -k^{\text{ex}}\|u\|^2 \tag{39}$$

since

$$(U_\Lambda[u]|p^e_\Lambda) = \sum_{x\in\mathcal{C}_\Lambda} U_\Lambda[u](x) = \sum_{\alpha\in\mathcal{A}} a_\alpha \sum_{x\in\mathcal{C}_\Lambda} p^e_\Lambda(x)\frac{u_\alpha(x_\alpha)}{p^e_\alpha(x_\alpha)} = \sum_{\alpha\in\mathcal{A}} a_\alpha \sum_{x_\alpha} u_\alpha(x_\alpha) = 0. \tag{40}$$

Finally, defining k as

$$k = -\max\left\{\frac{\langle u|\mathbb{T}_\Lambda u\rangle}{\langle u|u\rangle} : u \in \mathcal{V}_\Lambda \setminus \{0\}\right\} \tag{41}$$

we have by Eq. (39) $k \geq k^{\mathrm{ex}}$.

Now we can show that k, the opposite of the maximum eigenvalue of the operator $\mathbb{T}_\Lambda$, is the approximate relaxation rate. We have just seen that it is not smaller than the exact one, a result that can be intuitively understood by observing that the local equilibrium assumption implies that we are dealing with an evolution in a restricted probability space.

Proposition: Let $u^t = p^t - p^e$ be the solution of Eq. (35) with initial condition $u^0 = p^0 - p^e$. It vanishes exponentially with time or, more precisely, with k defined by Eq. (41), there exist v and $E^t \in \mathcal{V}_\Lambda$ such that

$$u^t = \mathrm{e}^{-kt}[v + E^t] \tag{42}$$

and

$$\lim_{t\to+\infty} E^t = 0. \tag{43}$$

Proof. We shall first find an exponential bound for $\|u^t\|$, by showing that there exist $t_0 > 0$ and $A > 0$ such that, for $t \geq t_0$, $u^t \in \mathcal{I}_0$ and

$$\|u^t\| \leq A\,\mathrm{e}^{-kt}. \tag{44}$$

Since $\lim_{t\to+\infty} u^t = 0$, given a positive constant b one can find $t_0 > 0$ such that $u^t \in \mathcal{I}_0$ and $\|u^t\| \leq b$ for $t \geq t_0$. Then, multiplying equation (35) by u^t and for $t \geq t_0$, we have

$$\begin{aligned} \frac{1}{2}\frac{d}{dt}\|u^t\|^2 &= \langle u^t|\mathbb{T}_\Lambda u^t\rangle + \langle u^t|R_\Lambda(u^t)\rangle \\ &\leq -k\|u^t\|^2 + M\|u^t\|^3 \\ &\leq -(k - Mb)\|u^t\|^2, \end{aligned} \tag{45}$$

which can be integrated to give

$$\|u^t\| \leq \mathrm{e}^{-(k-Mb)(t-t_0)}\|u^{t_0}\|. \tag{46}$$

Inserting the above result in the first inequality of Eq. (45) we obtain

$$\frac{1}{2}\frac{d}{dt}\|u^t\|^2 \leq -k\|u^t\|^2 + M\|u^{t_0}\|^3 \mathrm{e}^{-3(k-Mb)(t-t_0)}, \tag{47}$$

which can be integrated again to give

$$\|u^t\|^2 \leq \mathrm{e}^{-2k(t-t_0)}\|u^{t_0}\|^2 + \frac{M\|u^{t_0}\|^3}{k-3Mb}\left[\mathrm{e}^{-2k(t-t_0)} - \mathrm{e}^{-3(k-Mb)(t-t_0)}\right]. \tag{48}$$

The constant b can be chosen small enough to ensure that $k - 3Mb > 0$, and hence A in Eq. (44) can be given by

$$A = \mathrm{e}^{kt_0}\left[\|u^{t_0}\|^2 + \frac{M\|u^{t_0}\|^3}{k-3Mb}\right]^{1/2}. \tag{49}$$

Finally, since u^t can be expanded in eigenvectors of $\mathbb{T}_\Lambda$, our proposition is proven if we show that $\lim\limits_{t\to+\infty}\langle w|u^t\rangle \mathrm{e}^{kt}$ exists finite for any eigenvector w with $\|w\| = 1$. If this is the case v and E^t are given by $\lim\limits_{t\to+\infty} u^t\mathrm{e}^{kt}$ and $u^t\mathrm{e}^{kt} - v$ respectively. It can be noted that v has to be an eigenvector of $\mathbb{T}_\Lambda$ relative to $-k$.

Eq. (44) tells us that $\langle w|u^t\rangle\mathrm{e}^{kt}$ is bounded. If w corresponds to $-k$, let us consider an increasing sequence $\{t_n\}_{n\in\mathbb{N}}$ diverging to infinity such that t_0 is the value previously introduced. From Eq. (35) we have

$$\frac{d}{dt}\langle w|u^t\rangle = -k\langle w|u^t\rangle + \langle w|R_\Lambda(u^t)\rangle. \tag{50}$$

Using Eqs. (37) and (44) and integrating between t_n and t_{n+m} we obtain

$$\left|\langle w|u^{t_{n+m}}\rangle\mathrm{e}^{kt_{n+m}} - \langle w|u^{t_n}\rangle\mathrm{e}^{kt_n}\right| \leq \frac{MA^2}{k}\mathrm{e}^{-kt_n} \tag{51}$$

for all $m \in \mathbb{N}$, which shows that the sequence $\{\langle w|u^{t_n}\rangle\mathrm{e}^{kt_n}\}_{n\in\mathbb{N}}$ is a Cauchy one. Then the limit $\lim\limits_{n\to+\infty}\langle w|u^{t_n}\rangle\mathrm{e}^{kt_n}$ exists finite and thus, from the arbitrariety of the sequence $\{t_n\}_{n\in\mathbb{N}}$, also $\lim\limits_{t\to+\infty}\langle w|u^t\rangle\mathrm{e}^{kt}$ does it.

If w corresponds to $-\lambda < -k$ then

$$\frac{d}{dt}\langle w|u^t\rangle = -\lambda\langle w|u^t\rangle + \langle w|R_\Lambda(u^t)\rangle. \tag{52}$$

Using again Eqs. (37) and (44) and integrating, with manipulations similar to the previous ones, one can find two constants $B > 0$ and $C > 0$ such that, for $t \geq t_0$,

$$\left|\langle w|u^t\rangle\right| \leq B\,\mathrm{e}^{-2kt} + C\,\mathrm{e}^{-\lambda t}, \tag{53}$$

from which follows that $\lim\limits_{t\to+\infty}\langle w|u^t\rangle\mathrm{e}^{kt} = 0$.

IV. LOCAL EQUILIBRIUM KINETICS FOR THE WSME MODEL

After studying some features of the approximation, we are going to show its performance with an example, i.e. a particular, physically relevant, choice of the transition matrix. The goal is the reduction of the computational complexity of the kinetic problem from exponential to polynomial in the number of variables. We focus on a single bond-flip kinetics, that is we consider kinetics with only transitions between configurations that differ for no more than the state of one peptidic bond, and for which the detailed balance principle (28) is satisfied.

Given $x \in \mathcal{C}_\Lambda$ we denote by $\mu_k(x)$ the configuration obtained by x by turning the variable $x_{k,k}$, i.e. m_k, into $1 - x_{k,k}$. We have explicity

$$\mu_k(x)_{i,j} = \begin{cases} x_{i,j} \text{ if } j < k \text{ or } i > k; \\ x_{i,k-1}(1 - x_{k,k})x_{k+1,j} \text{ otherwise.} \end{cases} \tag{54}$$

A single bond-flip kinetics will be described by a transition matrix with the property $W_\Lambda(x \to x') = 0$ if $x' \notin \{x, \mu_1(x), \ldots, \mu_N\}$. For the matrix elements corresponding to the allowed transitions we set

$$W_\Lambda(x \to \mu_k(x)) = \nu(H_\Lambda(\mu_k(x)) - H_\Lambda(x)), \tag{55}$$

where $\nu : \mathbb{R} \to \mathbb{R}$ is a strictly positive function which satisfies detailed balance:

$$\nu(-\Delta) = \mathrm{e}^{\Delta}\nu(\Delta). \tag{56}$$

Condition (10) requires that

$$W_\Lambda(x \to x) = -\sum_{k=1}^{N} \nu(H_\Lambda(\mu_k(x)) - H_\Lambda(x)) \tag{57}$$

and it is easy to check that W_Λ is irreducible. The family of ν functions fulfilling Eq. (56) contains Metropolis and Glauber kinetic prescriptions, that are

$$\nu(\Delta) = \min\{1, \mathrm{e}^{-\Delta}\} \tag{58}$$

and

$$\nu(\Delta) = 1/(1 + \mathrm{e}^{\Delta}) \tag{59}$$

respectively.

To show the form of approximate kinetic equations we must explicitly write the r.h.s. of Eq. (15), i.e. the function f defined in (16). Writing probabilities as functions of the expectation values

$$\xi_{i,j} = \sum_{x \in \mathcal{C}_\Lambda} x_{i,j} P_\Lambda[p], \tag{60}$$

as anticipated in Sec. II, f takes the form

$$f_{i,j}(\xi) = \sum_{k=1}^{N} \sum_{x \in \mathcal{C}_\Lambda} (\mu_k(x)_{i,j} - x_{i,j}) \nu(H_\Lambda(\mu_k(x)) - H_\Lambda(x)) P_\Lambda[p](x) \tag{61}$$

where Eqs. (55), (57) and the substitution of x with $\mu_k(x)$ in the first term of the r.h.s. have been used. Since $\mu_k(x)_{i,j} = x_{i,j}$ if $k < i$ or $k > j$ we can also write

$$f_{i,j}(\xi) = \sum_{k=i}^{j} \sum_{x \in \mathcal{C}_\Lambda} (\mu_k(x)_{i,j} - x_{i,j}) \nu(H_\Lambda(\mu_k(x)) - H_\Lambda(x)) P_\Lambda[p](x). \tag{62}$$

In the following we will need the new variables $S_{i,j}$, which take value 1 if i and j are disordered peptide bond which delimit a string of consecutive ordered bonds. These variables are defined as

$$S_{i,j}(x) = (1 - m_i)(1 - m_j) \prod_{k=i+1}^{j-1} m_k = \begin{cases} 1 - x_{i,j} \text{ if } i = j; \\ (1 - x_{i,i})(1 - x_{j,j}) \text{ if } j = i + 1; \\ (1 - x_{i,i}) x_{i+1,j-1} (1 - x_{j,j}) \text{ if } 0 \leq i < j - 1 \leq N, \end{cases} \tag{63}$$

where we assume the boundary conditions $m_0 = m_{N+1} = 0$ which imply $x_{i,j} = 0$ if $i = 0$ or $j = N + 1$. It is not difficult to check that

$$x_{i,j} = \sum_{s=j}^{N} \sum_{r=1}^{i} S_{r-1,s+1}(x) \tag{64}$$

and

$$\sum_{j=k}^{N} \sum_{i=1}^{k} h_{i,j} x_{i,j} = \sum_{j=k}^{N} \sum_{i=1}^{k} \Delta_{i,j}^{k} S_{i-1,j+1}(x) \tag{65}$$

where, for $1 \leq i \leq k \leq j \leq N$, we have introduced the quantities

$$\Delta_{i,j}^{k} \doteq \sum_{s=k}^{j} \sum_{r=i}^{k} h_{r,s}. \tag{66}$$

Now we can return to Eq. (62). Notice that, for $i \leq k \leq j$,

$$\begin{aligned} & (\mu_k(x)_{i,j} - x_{i,j})\nu(H_\Lambda(\mu_k(x)) - H_\Lambda(x)) \\ = \; & (\mu_k(x)_{i,j} - x_{i,j})\nu\left(\sum_{s=k}^{N}\sum_{r=1}^{k} h_{r,s}(\mu_k(x)_{r,s} - x_{r,s})\right) \\ = \; & \mu_k(x)_{i,j}\nu\left(\sum_{s=k}^{N}\sum_{r=1}^{k} h_{r,s}\mu_k(x)_{r,s}\right) - x_{i,j}\nu\left(-\sum_{s=k}^{N}\sum_{r=1}^{k} h_{r,s}x_{r,s}\right). \end{aligned} \tag{67}$$

The first equality follows by the relation $\mu_k(x)_{i,j} = x_{i,j}$ for $i > k$ or $j < k$. The second one by the fact that if $i \leq k \leq j$ and $x_{i,j} = 1$ then $\mu_k(x)_{r,s} = 0$ for $r \leq k$ and $s \geq k$, and vice–versa.

We will concentrate only on the term

$$x_{i,j}\nu\left(-\sum_{s=k}^{N}\sum_{r=1}^{k} h_{r,s}x_{r,s}\right) \tag{68}$$

since we can obtain the first one by replacing $h_{i,j}$ with $-h_{i,j}$ and x with $\mu_k(x)$. Eq. (65) allows us to write

$$x_{i,j}\nu\left(-\sum_{s=k}^{N}\sum_{r=1}^{k} h_{r,s}x_{r,s}\right) = x_{i,j}\nu\left(-\sum_{s=k}^{N}\sum_{r=1}^{k} \Delta^k_{r,s}S_{r-1,s+1}(x)\right). \tag{69}$$

The last equality is nontrivial only for configurations $x \in \mathcal{C}_\Lambda$ such that $x_{i,j} = 1$, and hence $x_{k,k} = 1$. We now observe that, if $x_{k,k} = 1$, varing i in $\{1, \ldots, k\}$ and j in $\{k, \ldots, N\}$ one can find one and only one couple of indices (r, s) such that $S_{r-1,s+1}(x) = 1$. This couple of indices delimit the string of adjacent ordered bonds containing k. It follows that

$$x_{i,j}\nu\left(-\sum_{s=k}^{N}\sum_{r=1}^{k} h_{r,s}x_{r,s}\right) = x_{i,j}\sum_{s=k}^{N}\sum_{r=1}^{k} \nu(-\Delta^k_{r,s})S_{r-1,s+1}(x). \tag{70}$$

Finally, observing that for $i \leq k \leq j$ and $r \leq k \leq s$

$$x_{i,j}S_{r-1,s+1}(x) = \begin{cases} 0 \text{ if } r > i \text{ or } s < j; \\ S_{r-1,s+1}(x) \text{ otherwise,} \end{cases} \tag{71}$$

we reach the result

$$x_{i,j}\nu\left(-\sum_{s=k}^{N}\sum_{r=1}^{k} h_{r,s}x_{r,s}\right) = \sum_{s=j}^{N}\sum_{r=1}^{i} \nu(-\Delta^k_{r,s})S_{r-1,s+1}(x) \tag{72}$$

that makes possible to rewrite f in a simpler way.

Substituting Eq. (72) in Eq. (67) and using Eq. (62) we obtain

$$\begin{aligned} f_{i,j}(\xi) &= \sum_{k=i}^{j}\sum_{s=j}^{N}\sum_{r=1}^{i}\nu(\Delta^k_{r,s})\sum_{x\in\mathcal{C}_\Lambda} S_{r-1,s+1}(\mu_k(x))P_\Lambda[p](x) + \\ &- \sum_{k=i}^{j}\sum_{s=j}^{N}\sum_{r=1}^{i}\nu(-\Delta^k_{r,s})\sum_{x\in\mathcal{C}_\Lambda} S_{r-1,s+1}(x)P_\Lambda[p](x) \end{aligned} \tag{73}$$

that contains expectation values of two kinds of observables. The sum

$$\sum_{x\in\mathcal{C}_\Lambda} S_{r-1,s+1}(x)P_\Lambda[p](x) \tag{74}$$

is easily computable since it is the expectation value of a linear combination of $x_{i,j}$'s. With a slight abuse of notation we will still denote this sum with $S_{r-1,s+1}(\xi)$. One can verify that

$$S_{i,j}(\xi) = \begin{cases} 1-\xi_{i,j} \text{ if } i=j; \\ \xi_{i,j}-\xi_{i,i}-\xi_{j,j}+1 \text{ if } j=i+1; \\ \xi_{i,j}-\xi_{i,j-1}-\xi_{i+1,j}+\xi_{i+1,j-1} \text{ if } 0\le i<j-1\le N, \end{cases} \tag{75}$$

where we assume $\xi_{i,j}=0$ if $i=0$ or $j=N+1$.

The computation of

$$\sum_{x\in\mathcal{C}_\Lambda} S_{r-1,s+1}(\mu_k(x))P_\Lambda[p](x) \tag{76}$$

is slightly more involved. Notice first that for $r\le k\le s$

$$S_{r-1,s+1}(\mu_k(x)) = S_{r-1,k}(x)S_{k,s+1}(x), \tag{77}$$

which vanishes if $x_{k,k}=m_k=1$. Let us then define the clusters

$$\Lambda^d_k \doteq \{(i,j)\in\Lambda : j\le k\}, \tag{78}$$

$$\Lambda^u_k \doteq \{(i,j)\in\Lambda : j\ge k\}, \tag{79}$$

$$\mathcal{R}_k \doteq \{(i,j)\in\Lambda : j=k\}, \tag{80}$$

depicted in Fig. 1, and introduce their probability distributions

$$p_{\Lambda^d_k}(x_{\Lambda^d_k}) \doteq \sum_{x_{\Lambda\setminus\Lambda^d_k}} P_\Lambda[p](x), \tag{81}$$

$$p_{\Lambda^u_k}(x_{\Lambda^u_k}) \doteq \sum_{x_{\Lambda\setminus\Lambda^u_k}} P_\Lambda[p](x), \tag{82}$$

$$p_{\mathcal{R}_k}(x_{\mathcal{R}_k}) \doteq \sum_{x_{\Lambda\setminus\mathcal{R}_k}} P_\Lambda[p](x). \tag{83}$$



FIG. 1: Graphical representation of the clusters Λ_k^d, Λ_k^u and $\mathcal{R}_k$

Exploiting the constraints in the same way as in [9] it is not difficult to obtain the relations

$$P_\Lambda[p](x) = \frac{p_{\Lambda_k^d}(x_{\Lambda_k^d}) p_{\Lambda_k^u}(x_{\Lambda_k^u})}{p_{\mathcal{R}_k}(x_{\mathcal{R}_k})} \tag{84}$$

and

$$p_{\mathcal{R}_k}(0, \ldots, 0) = p_{(k,k)}(0) = 1 - \xi_{k,k}. \tag{85}$$

The quantity in (76) can then be rewritten as

$$\begin{aligned} & \sum_{x \in \mathcal{C}_\Lambda} S_{r-1,s+1}(\mu_k(x)) P_\Lambda[p](x) \\ = & \sum_{x \in \mathcal{C}_\Lambda} S_{r-1,k}(x) S_{k,s+1}(x) P_\Lambda[p](x) \\ = & \sum_{x \in \mathcal{C}_\Lambda} S_{r-1,k}(x) S_{k,s+1}(x) \frac{p_{\Lambda_k^d}(x_{\Lambda_k^d}) p_{\Lambda_k^u}(x_{\Lambda_k^u})}{p_{\mathcal{R}_k}(x_{\mathcal{R}_k})} \\ = & \frac{1}{1 - \xi_{k,k}} \sum_{x \in \mathcal{C}_\Lambda} S_{r-1,k}(x) S_{k,s+1}(x) p_{\Lambda_k^d}(x_{\Lambda_k^d}) p_{\Lambda_k^u}(x_{\Lambda_k^u}), \end{aligned} \tag{86}$$

since the only configurations which contribute to the sum are those with $x_{k,k} = 0$. In addition, due to the vanishing of $x_{k,k}$, the above sum can be factored as follows:

$$\begin{aligned} & \sum_{x \in \mathcal{C}_\Lambda} S_{r-1,s+1}(\mu_k(x)) P_\Lambda[p](x) \\ = & \frac{1}{1 - \xi_{k,k}} \sum_{x_{\Lambda_k^d}} S_{r-1,k}(x_{\Lambda_k^d}) p_{\Lambda_k^d}(x_{\Lambda_k^d}) \sum_{x_{\Lambda_k^u}} S_{k,s+1}(x_{\Lambda_k^u}) p_{\Lambda_k^u}(x_{\Lambda_k^u}) \end{aligned}$$

$$
\begin{aligned}
&= \frac{1}{1-\xi_{k,k}} \sum_{x\in\mathcal{C}_\Lambda} S_{r-1,k}(x) P_\Lambda[p](x) \sum_{x\in\mathcal{C}_\Lambda} S_{k,s+1}(x) P_\Lambda[p](x) \\
&= \frac{S_{r-1,k}(\xi) S_{k,s+1}(\xi)}{1-\xi_{k,k}}. \qquad (87)
\end{aligned}
$$

Plugging Eq. (87) into Eq. (73) we find for f the final form

$$
f_{i,j}(\xi) = \sum_{k=i}^{j} \sum_{s=j}^{N} \sum_{r=1}^{i} \left[\nu(\Delta^k_{r,s}) \frac{S_{r-1,k}(\xi) S_{k,s+1}(\xi)}{S_{k,k}(\xi)} - \nu(-\Delta^k_{r,s}) S_{r-1,s+1}(\xi) \right]. \qquad (88)
$$

¿From the above results one sees that the computational complexity of the r.h.s. of Eq. (15) is polynomial in the number of variables, and precisely of order N^5. Moreover it must be mentioned that writing directly an equation for the variables $S_{i,j}$ the computational complexity lowers to the order N^3, which is useful in practical computations.

V. EQUIVALENCE BETWEEN LOCAL EQUILIBRIUM APPROACH AND PATH PROBABILITY METHOD

We conclude our paper by showing that for the WSME model there are no differences between local equilibrium approach and path probability method (PPM) [24, 25, 26], a variational approximate technique for the study of out of equilibrium systems that generalzes the CVM to the kinetics.

The PPM has a natural application to discrete time evolution, hence it is useful to turn to this one and then to recover the continuous case with a limiting process. We say that $T_\Lambda(x' \to x) \geq 0$ is a transition probability from the configuration x' to x if

$$
\sum_{x\in\mathcal{C}_\Lambda} T_\Lambda(x' \to x) = 1. \qquad (89)
$$

If W_Λ is the transition probability per unit time as listed in Eq. (9), let T^τ_Λ be a transition probability such that

$$
T^\tau_\Lambda(x' \to x) = \delta_{x,x'} + \tau W_\Lambda(x' \to x) + \tau^2 E^\tau_\Lambda(x' \to x) \qquad (90)
$$

where E^τ_Λ is a bounded function of τ when $\tau \to 0$ and δ the Kronecker symbol. The discrete version of Eq. (9) then reads

$$
p^{t+\tau}_\Lambda(x) = \sum_{x'\in\mathcal{C}_\Lambda} T^\tau_\Lambda(x' \to x) p^t_\Lambda(x') \qquad (91)
$$

for all $t \in \{n\tau\}_{n\in\mathbb{N}}$ and, as a consequence of Eq. (90), given $t \in \mathbb{R}_+$, in the limit $\tau \to 0$ and $n \to \infty$ such that $n\tau \to t$, the solution of this equation reduces to that of Eq. (9) with the same initial condition.

In PPM [24, 25, 26] one assumes that T^τ_Λ has the form

$$T^\tau_\Lambda(x' \to x) = \Theta^\tau_\Lambda(x') \prod_{\alpha\in\mathcal{A}} \left[T^\tau_\alpha(x'_\alpha \to x_\alpha)\right]^{a_\alpha}, \tag{92}$$

where Θ^τ_Λ ensures normalization (89) and $\mathcal{A}$ and a_α are the same of Sec. II.

Let $\mathcal{D}^2_\Lambda$ be the set of functions $q = \{q_\alpha\}_{\alpha\in\mathcal{A}}$ such that $q_\alpha : \mathcal{C}_\alpha \times \mathcal{C}_\alpha \to \mathbb{R}$ is not negative and $\sum_{x'_{\alpha\setminus\beta}} \sum_{x_{\alpha\setminus\beta}} q_\alpha(x'_\alpha, x_\alpha) = q_\beta(x'_\beta, x_\beta)$ for $\beta \subset \alpha$. For $q \in \mathcal{D}^2_\Lambda$ define then the function $Q_\Lambda[q] : \mathcal{C}_\Lambda \times \mathcal{C}_\Lambda \to \mathbb{R}_+$ as

$$Q_\Lambda[q](x', x) = \prod_{\alpha\in\mathcal{A}} \left[q_\alpha(x'_\alpha, x_\alpha)\right]^{a_\alpha}. \tag{93}$$

It can be checked that q_α is the marginal of $Q_\Lambda[q]$ relative to the cluster α, i.e.

$$q_\alpha(x'_\alpha, x_\alpha) = \sum_{x'_{\Lambda\setminus\alpha}} \sum_{x_{\Lambda\setminus\alpha}} Q_\Lambda[q](x', x). \tag{94}$$

The quantity $T^\tau_\Lambda(x' \to x)p^t_\Lambda(x')$, where $p^t_\Lambda(x')$ is the solution of Eq. (91), has a physical meaning: it is the joint probability of finding the system in the state x' at time t and in the state x at time $t+\tau$. The idea of the PPM is to replace this probability with an approximate one, which factors as in Eq. (93), i.e. to replace $T^\tau_\Lambda(x' \to x)p^t_\Lambda(x')$ with $Q_\Lambda[q^{t,\tau}](x', x)$, and then to compute the marginal distribution at the time $t + \tau$ as

$$p^{t+\tau}_\alpha(x_\alpha) = \sum_{x'\in\mathcal{C}_\Lambda} \sum_{x_{\Lambda\setminus\alpha}} Q_\Lambda[q^{t,\tau}](x', x) = \sum_{x'_\alpha} q^{t,\tau}_\alpha(x'_\alpha, x_\alpha). \tag{95}$$

$q^{t,\tau}_\alpha$ is then an approximation of the above joint distribution relative to the cluster α. The function $q^{t,\tau}$ must contain information about the state at time t and the kinetic prescription. As a consequence of the relation

$$\sum_{x'_{\Lambda\setminus\alpha}} \sum_{x\in\mathcal{C}_\Lambda} T^\tau_\Lambda(x' \to x)p^t_\Lambda(x') = p^t_\alpha(x'_\alpha), \tag{96}$$

we impose to $q^{t,\tau}$ the constraints

$$\sum_{x'_{\Lambda\setminus\alpha}} \sum_{x\in\mathcal{C}_\Lambda} Q_\Lambda[q^{t,\tau}](x', x) = \sum_{x_\alpha} q^{t,\tau}_\alpha(x'_\alpha, x_\alpha) = p^t_\alpha(x'_\alpha), \tag{97}$$

that it has to be satisfied for all $\alpha \in \mathcal{A}$.

The kinetic generalization of the Kikuchi free energy F_Λ is

$$K_\Lambda^\tau[q] = \sum_{\alpha\in\mathcal{A}} a_\alpha \sum_{x'_\alpha}\sum_{x_\alpha} \left[\ln q_\alpha(x'_\alpha, x_\alpha) - \ln T_\alpha^\tau(x'_\alpha \to x_\alpha)\right] q_\alpha(x'_\alpha, x_\alpha), \tag{98}$$

and $q^{t,\tau}$ is chosen as the distribution $q \in \mathcal{D}_\Lambda^2$ which minimizes K_Λ^τ with the constraints (97). If the factorization assumption were exact, this choice would give the exact marginals of the joint probability $T_\Lambda^\tau(x' \to x)p_\Lambda^t(x')$.

It can be checked that conditions $\lim_{\tau\to 0} q_\alpha^{t,\tau}(x'_\alpha, x_\alpha) = \delta_{x_\alpha, x'_\alpha} p_\alpha^t(x_\alpha)$ hold for all $\alpha \in \mathcal{A}$. These allow us to summarize PPM, for continuous time evolution, as follows:

$$\begin{cases} \displaystyle\frac{d}{dt}p_\alpha^t(x_\alpha) = \frac{\partial}{\partial\tau}\sum_{x'_\alpha} q_\alpha^{t,\tau}(x'_\alpha, x_\alpha)\bigg|_{\tau=0} \\ q^{t,\tau} = \operatorname{minarg}\{K_\Lambda^\tau[q] : q \in \mathcal{D}_\Lambda^2[p^t]\}, \end{cases} \tag{99}$$

where, for $p \in \mathcal{D}_\Lambda$, $\mathcal{D}_\Lambda^2[p]$ denotes the set of $q \in \mathcal{D}_\Lambda^2$ such that $\sum_{x_\alpha} q_\alpha(x'_\alpha, x_\alpha) = p_\alpha(x_\alpha)$. In the following we shall prove the equivalence between the local equilibrium approach and PPM by showing that the problem (99) leads to Eq. (12). For this purpose we need the following proposition.

Proposition: Let q^τ be the minimum of the function K_Λ^τ in $\mathcal{D}_\Lambda^2[p]$. Then a function $R_\Lambda^\tau : \mathcal{C}_\Lambda \to \mathbb{R}_+$ exists such that

$$Q_\Lambda[q^\tau](x', x) = R_\Lambda^\tau(x')T_\Lambda^\tau(x' \to x)P_\Lambda[p](x'). \tag{100}$$

Moreover R_Λ^τ can be written as

$$R_\Lambda^\tau(x) = 1 + \tau r_\Lambda^\tau(x) \tag{101}$$

where $r_\Lambda^\tau : \mathcal{C}_\Lambda \to \mathbb{R}$ is a bounded function of τ, when τ approaches zero, with the property

$$\sum_{x_{\Lambda\setminus\alpha}} r_\Lambda^\tau(x)P_\Lambda[p](x) = 0 \tag{102}$$

for all $\alpha \in \mathcal{A}$.

Before proving the proposition, we show how it leads to the equivalence of the two methods. From now on we denote by q^τ the minimum of K_Λ^τ in $\mathcal{D}_\Lambda^2[p]$. We want to show that

$$\frac{\partial}{\partial\tau}\sum_{x'_\alpha} q_\alpha^\tau(x'_\alpha, x_\alpha)\bigg|_{\tau=0} = \sum_{x'\in\mathcal{C}_\Lambda} W_\alpha(x' \to x_\alpha)P_\Lambda[p](x'), \tag{103}$$

from which it follows that the PPM problem (99) coincides with the local equilibrium approach, Eq. (12). Eqs. (100), (101) and (90) allow us to obtain

$$\begin{aligned} Q_\Lambda[q^\tau](x',x) &= R^\tau_\Lambda(x')T^\tau_\Lambda(x'\to x)P_\Lambda[p](x') \\ &= \left[\delta_{x,x'}+\tau\delta_{x,x'}r^\tau_\Lambda(x')+\tau W_\Lambda(x'\to x)+\tau^2E^\tau_\Lambda(x'\to x)\right]P_\Lambda[p](x'), \end{aligned} \tag{104}$$

where, as before, E^τ_Λ is a bounded function of τ when τ approaches zero. Then, summing and using Eq. (102), we have

$$\begin{aligned} \sum_{x'_\alpha} q^\tau_\alpha(x'_\alpha,x_\alpha) &= \sum_{x_{\Lambda\setminus\alpha}}\sum_{x'\in\mathcal{C}_\Lambda} Q_\Lambda[q^\tau](x',x) \\ &= p_\alpha(x_\alpha)+\tau\sum_{x_{\Lambda\setminus\alpha}} r^\tau_\Lambda(x)P_\Lambda[p](x)+\tau\sum_{x_{\Lambda\setminus\alpha}}\sum_{x'\in\mathcal{C}_\Lambda} W_\Lambda(x'\to x)P_\Lambda[p](x') \\ &+ \tau^2\sum_{x_{\Lambda\setminus\alpha}}\sum_{x'\in\mathcal{C}_\Lambda} E^\tau_\Lambda(x'\to x) \\ &= p_\alpha(x_\alpha)+\tau\sum_{x_{\Lambda\setminus\alpha}}\sum_{x'\in\mathcal{C}_\Lambda} W_\Lambda(x'\to x)P_\Lambda[p](x') \\ &+ \tau^2\sum_{x_{\Lambda\setminus\alpha}}\sum_{x'\in\mathcal{C}_\Lambda} E^\tau_\Lambda(x'\to x), \end{aligned} \tag{105}$$

that, remembering the definition (13) of $W_\alpha(x'\to x_\alpha)$, gives the expected result (103). Let us move now to the proof of the proposition.

Proof. Let us denote by $\mathcal{A}_M$ the family of clusters, the maximal clusters, that are not enclosed in any others and by $\mathcal{A}_m$ those which are intersections of at least two maximal clusters. Notice that $\mathcal{A}_M\cap\mathcal{A}_m=\emptyset$ and $\mathcal{A}_M\cup\mathcal{A}_m=\mathcal{A}$. In the case of the WSME model $\mathcal{A}_M$ is the set of all square plaquettes and the triangles lying on the diagonal boundary, while $\mathcal{A}_m$ is the set of internal nearest–neighbour pairs and single nodes.

In addition, given $\beta\in\mathcal{A}_m$, let us introduce the set

$$\mathcal{A}^\beta_M \doteq \{\alpha\in\mathcal{A}_M:\beta\subset\alpha\}. \tag{106}$$

In the same way, for $\alpha\in\mathcal{A}_M$,

$$\mathcal{A}^\alpha_m \doteq \{\beta\in\mathcal{A}_m:\beta\subset\alpha\}. \tag{107}$$

The minimization of K^τ_Λ in $\mathcal{D}_\Lambda[p]$ is addressed by the method of Lagrange multiplyers. We consider the function

$$K^\tau_\Lambda[q;\lambda] \doteq K^\tau_\Lambda[q]-\sum_{\alpha\in\mathcal{A}}\sum_{x'_\alpha}\lambda_\alpha(x'_\alpha)\left[\sum_{x_\alpha} q_\alpha(x'_\alpha,x_\alpha)-p_\alpha(x'_\alpha)\right]$$

$$+\sum_{\beta\in\mathcal{A}_m}\sum_{\alpha\in\mathcal{A}_M^\beta}\sum_{x_\beta}\sum_{x'_\beta}\lambda_{\alpha\to\beta}(x_\beta,x'_\beta)\left[\sum_{x_{\alpha\setminus\beta}}\sum_{x'_{\alpha\setminus\beta}}q_\alpha(x'_\alpha,x_\alpha)-q_\beta(x'_\beta,x_\beta)\right]. \quad (108)$$

Taking derivatives with respect to the variables $q_\alpha(x'_\alpha,x_\alpha)$ we obtain the equations

$$a_\alpha\ln q^\tau_\alpha(x'_\alpha,x_\alpha)-a_\alpha\ln T^\tau_\alpha(x'_\alpha\to x_\alpha)+\sum_{\beta\in\mathcal{A}_m^\alpha}\lambda^\tau_{\alpha\to\beta}(x_\beta,x'_\beta)-\lambda^\tau_\alpha(x'_\alpha)+a_\alpha=0 \quad (109)$$

and

$$a_\beta\ln q^\tau_\beta(x'_\beta,x_\beta)-a_\beta\ln T^\tau_\beta(x'_\beta\to x_\beta)-\sum_{\alpha\in\mathcal{A}_M^\beta}\lambda^\tau_{\alpha\to\beta}(x_\beta,x'_\beta)-\lambda^\tau_\beta(x'_\beta)+a_\beta=0, \quad (110)$$

that are satisfied by q^τ for $\alpha\in\mathcal{A}_M$ and $\beta\in\mathcal{A}_m$. Summing now the first over $\mathcal{A}_M$, the second ones over $\mathcal{A}_m$ and the results together, we arrive at Eq. (100):

$$\begin{aligned}Q_\Lambda[q^\tau](x',x) &= \prod_{\alpha\in\mathcal{A}}\mathrm{e}^{\lambda^\tau_\alpha(x'_\alpha)-a_\alpha}\prod_{\alpha\in\mathcal{A}}\left[T^\tau_\alpha(x'_\alpha\to x_\alpha)\right]^{a_\alpha}\\ &= R^\tau_\Lambda(x')T^\tau_\Lambda(x'\to x)P_\Lambda[p](x') \qquad (111)\end{aligned}$$

with a suitable function R^τ_Λ.

The relations $\lim_{\tau\to0}q^\tau_\alpha(x'_\alpha,x_\alpha)=\delta_{x_\alpha,x'_\alpha}p_\alpha(x_\alpha)$ and $\lim_{\tau\to0}T^\tau_\Lambda(x'\to x)=\delta_{x,x'}$ imply then

$$\lim_{\tau\to0}R^\tau_\Lambda(x)=1 \quad (112)$$

and from Eq. (111) it follows that a bounded function $r^\tau_\Lambda:\mathcal{C}_\Lambda\to\mathbb{R}$ exists such that Eq. (101) holds.

We conclude the proof by showing the validity of Eq. (102). Using Eq. (89) we find

$$\sum_{x\in\mathcal{C}_\Lambda}Q_\Lambda[q^\tau](x',x)=R^\tau_\Lambda(x')P_\Lambda[p](x'), \quad (113)$$

and using Eq. (101) we obtain

$$\begin{aligned}p_\alpha(x') &= \sum_{x_\alpha}q^\tau_\alpha(x'_\alpha,x_\alpha)=\sum_{x'_{\Lambda\setminus\alpha}}\left[\sum_{x\in\mathcal{C}_\Lambda}Q_\Lambda[q^\tau](x',x)\right]\\ &= \sum_{x'_{\Lambda\setminus\alpha}}R^\tau_\Lambda(x')P_\Lambda[p](x')\\ &= \sum_{x'_{\Lambda\setminus\alpha}}P_\Lambda[p](x')+\tau\sum_{x'_{\Lambda\setminus\alpha}}r^\tau_\Lambda(x')P_\Lambda[p](x')\\ &= p_\alpha(x')+\tau\sum_{x'_{\Lambda\setminus\alpha}}r^\tau_\Lambda(x')P_\Lambda[p](x'), \qquad (114)\end{aligned}$$

from which Eq. (102) follows.

VI. CONCLUSIONS

We have given proofs of a few rigorous results we have announced in [1], concerning the properties of the local equilibrium approach to the WSME model. In particular, we have proven that (i) the free energy decreases with time, (ii) the exact equilibrium is recovered in the infinite time limit, (iii) the equilibration rate is an upper bound of the exact one, (iv) computational complexity is polynomial in the number of variables, and (v) the local equilibrium approach is equivalent to the path probability method. We have also reported the detailed form of the kinetic equations.

It is important here to stress that our proofs depend only on the assumption of a factorization property for the probability distribution of the model, and hence can be easily carried over to other models with the same property.

Our results also shed new light on the physical meaning of the path probability method.

---

[1] M. Zamparo and A. Pelizzola, Phys. Rev. Lett. **97**, 068106 (2006).

[2] H. Wako and N. Saitô, J. Phys. Soc. Jpn **44**, 1931 (1978).

[3] H. Wako and N. Saitô, J. Phys. Soc. Jpn **44**, 1939 (1978).

[4] V. Muñoz, P.A. Thompson, J. Hofrichter and W.A. Eaton, Nature **390**, 196 (1997).

[5] V. Muñoz, E.R. Henry, J. Hofrichter and W.A. Eaton, Proc. Natl. Acad. Sci. U.S.A. **95**, 5872 (1998).

[6] V. Muñoz and W.A. Eaton, Proc. Natl. Acad. Sci. U.S.A. **96**, 11311 (1999).

[7] A. Flammini, J.R. Banavar and A. Maritan, Europhys. Lett. **58**, 623 (2002).

[8] P. Bruscolini and A. Pelizzola, Phys. Rev. Lett. **88**, 258101 (2002).

[9] A. Pelizzola, J. Stat. Mech., P11010 (2005).

[10] R. Kikuchi, Phys. Rev. **81**, 988 (1951).

[11] G. An, J. Stat. Phys. **52**, 727 (1988).

[12] A. Pelizzola, J. Phys. A **38**, R309 (2005).

[13] K. Itoh and M. Sasai, Proc. Natl. Acad. Sci. U.S.A. **101**, 14736 (2004).

[14] K. Itoh and M. Sasai, Chem. Phys. **307**, 121 (2004).

[15] E.R. Henry, W.A. Eaton, Chem. Phys. **307**, 163 (2004).

[16] K. Itoh and M. Sasai, Proc. Natl. Acad. Sci. U.S.A. **103**, 7298 (2006).

[17] H. Abe and H. Wako, Phys. Rev. E **74**, 011913 (2006).

[18] V.I. Tokar and H. Dreyssé, Phys. Rev. E **68**, 011601 (2003).

[19] V.I. Tokar and H. Dreyssé, J. Phys. Cond. Matter **16**, S2203 (2004).

[20] V.I. Tokar and H. Dreyssé, Phys. Rev. E **71**, 031604 (2005).

[21] K. Kawasaki, in *Phase Transitions and Critical Phenomena*, vol. 2, ed. C. Domb and M.S. Green (Academic, London, 1972).

[22] J.-F. Gouyet, M. Plapp, W. Dieterich and P. Maass, Adv. Phys. **52**, 523 (2003).

[23] T. Petermann e P. De Los Rios, J. Theor. Biol. **229**, 1 (2004).

[24] T. Ishii, Progr. Theor. Phys. Suppl. **115**, 243 (1994).

[25] F. Ducastelle, Progr. Theor. Phys. Suppl. **115**, 255 (1994).

[26] K. Wada e M. Kaburagi, Progr. Theor. Phys. Suppl. **115**, 273 (1994).

[27] N.G. Van Kampen, *Stochastic processes in physics and chemistry* (North–Holland, Amsterdam, 1981).

[28] J.L. Willems, *Stability theory of dynamical systems* (Nelson, London, 1970).